Conceptos generales de la IMRT rotatoria. Diferentes alternativas (Tomotherapy - VMAT - RapidArc). Consideraciones desde el punto de vista físico
William Parker

Centre universitaire
de santé McGill

McGill University
Health Centre

Research
Institute

Cancer
Centre

MCI/RVH
Adult

MCH
Children's

Shriners

Radiation Oncology - McGill

- Instalaciones de 70,000 pies cuadrados - 2015
- 150 personal incluyen:
 - 14 MDs + 14 residentes
 - 14 físicos + 9 dosimetristas + 4 res
 - 45 tecnólogos
- 3000 nuevos pacientes por año
- 160 tratamientos diarios
- 60 MD diarios
- 7 aceleradores – 6 TB + 1 CK
- 3 CTs
- 1 braquiterapia
- 1 MRI
- ARIA - Eclipse

D S1 RADIATION ONCOLOGY / MD CONSULTATION

© copyright 8 May 2015 McGill University Health Centre.

Conceptos generales de la IMRT rotatoria. Diferentes alternativas (Tomotherapy - VMAT - RapidArc). Consideraciones desde el punto de vista físico
William Parker

IMRT definicion

- Intensity Modulated Radiation Therapy
- "La IMRT se basa en el uso de intensidades de haz de radiación no uniformes optimizadas que inciden en el paciente" (NIH CWG, IJROBP, 2001).
- Gran número de campos
- Alto número de MUs (5 - 10 veces más que la CRT 3D)
- Largo tiempo de tratamiento

VMAT definición

- Volumetric modulated arc therapy
- IMRT realizado en un solo arco (Otto 2008)

Tiempo, esfuerzo, costo vs. beneficio

Year	Investigators	Milestone
1982	Brahme et al. [2]	Solving of an 'inverse' problem of rotational beam fluence to deliver a uniform dose to a donut-shaped target
1984	Brahme [15]	First commercial MLCs patented by Brahme and commercialized by Scanditronix
1988	Brahme [8]	First paper on algebraic inverse planning
1989	Webb [9]	Cast the inverse planning problem as an optimization problem that minimizes an objective or 'cost' function
1990	Bortfeld et al. [10]	Developed algebraic/iterative optimization for inverse planning
1991	Boyer et al. [48]	Developed principle of multiple segmented-field IMRT
1992	Convey and Rosenbloom [18]	Developed principle of sliding-window technique IMRT
1992	Carol [16]	First full IMRT system, based on serial tomotherapy, was introduced. NOMOS MIMiC binary MLC and simulated annealing optimization inverse planning (Peacock Plan)
1993	Mackie et al. [17]	Concept of helical tomotherapy
1994	Svensson et al. [19]	Independent finding of optimal and analytic solutions for sliding-window leaf trajectory problem by three research groups (Karolinska group in Stockholm, MSKCC group in New York, and DKFZ group in Heidelberg)
	Spirou and Chui [20]	
	Stein et al. [21]	
1994	Bortfeld et al. [49]	First multiple-static-field (MSF) experiments
1994	Carol et al. [50]	First IMRT treatment using serial tomotherapy (NOMOS MIMiC)
1995	Ling et al. [11]	First MLC-based IMRT at the Memorial Sloan Kettering Cancer Center
1995	Yu [28]	Intensity-modulated arc therapy; proposed as a tomotherapy-mimicking IMRT using a linac
2002	-	Tomotherapy released commercially
2007	Otto [26]	Volumetric-modulated arc therapy: IMRT in a single gantry arc
2010	Bogdanich [38]	New York Times IMRT accident reports

Peacock™: A System for Planning and Rotational Delivery of Intensity-Modulated Fields

Mark P. Carol

Nomos® Corporation, 2591 Wexford-Bayne

1992

I. INTRODUCTION

The effectiveness of radiation therapy in the clinical practice is limited by the oncologist's ability to

Helical Tomotherapy

1993

Mackie et al. Med Phys. 20(6) 1993

Experiencia McGill con Tomotherapy

- 2007-2014
- Mas de 1,500 pacientes
- Todos sitios anatómicos
- Primera maquina de IGRT y rotatoria
- TPS lento, pero distribuciones exquisitas
- Campos largos
- Tecnólogos gustaban mucho
- “single point setup”

Craniospinal radiotherapy

Pediatric CSI

PF dose reduction

Boost to mets

Aseguramiento de calidad integrado

Planificación adaptable de la IMRT

- Calcular la dosis en cada fracción
- "dosis deformada" para volver a la tomografía computarizada de planificación o al último conjunto de imágenes
- Calcular la DVH y las estadísticas de la dosis acumulada

Tomotherapy

- Tiempos de planificación largos
- Tiempo de toma de imágenes ~ 5 minutos por MVCT
- Tiempo de tratamiento ~ 10 - 15 minutos de haz
- 15 pacientes por día máximo

VMAT concepto

Intensity-modulated arc therapy with dynamic multileaf collimation: an alternative to tomotherapy

Cedric X Yu
William Beaumont Hospital, Royal Oak, MI, USA

1995

Received 9 February 1995, in final form 20 April 1995

IMAT with dynamic multileaf collimation

1447

(a)

VMAT solución práctica

2007

Volumetric modulated arc therapy: IMRT in a single gantry arc

Karl Otto^{B)}

Vancouver Cancer Centre, BC Cancer Agency, Vancouver, British Columbia V5Z 4E6, Canada

(Received 25 June 2007; revised 21 September 2007; accepted for publication 5 November 2007; published 26 December 2007)

En particular, la técnica VMAT está diseñada para que se puedan entregar planos optimizados:

- Eficientemente en un solo arco
- Con alta conformidad de dosis usando un rango de 360 grados de direcciones de pórtico
- Con precisión con muestreo de alta resolución de las direcciones de los haces durante la planificación

Otto. Med Phys. 2008.

¿Por qué rotar?

- Mejores planes de tratamiento
 - El aumento de los grados de libertad resulta en mejores planes de tratamiento
- Entrega eficiente
 - Sin arranques ni paradas
 - Menos unidades de monitor
 - Tratamiento más corto

Linac con OBI - compite bien con la Tomoterapia

Planificación de la IMRT - # de campos

5 Fields

7 Fields

9 Fields

La adición de más campos proporciona más grados de libertad para la optimización

Lo que ofrece una mayor posibilidad de adaptar la dosis al PTV y a los OAR.

Planificación de la IMRT - # de campos

Fields	Dose Prescription	Field Alignments	Plan Objectives	Optimization Objectives	Dose Statistics	Calculation Models	Plan Sum					
View	DVH Line	Structure	Approval Status	Plan	Course	Volume [cm ³]	Dose Cover. [%]	Sampling Cover. [%]	Min Dose [Gy]	Max Dose [Gy]	Mean Dose [Gy]	
☒		BODY	Unapproved	IMRT-5F-Ant	C1 Num of fields	21322.1	100.0	98.5	0.683	79.857	21.729	
☒		BODY	Unapproved	IMRT-7F	C1 Num of fields	21322.1	100.0	98.5	0.604	76.276	19.742	
☒		BODY	Unapproved	IMRT-9F	C1 Num of fields	21322.1	100.0	98.5	0.657	76.099	19.805	

IMRT - Ángulos de campo

7F Anterior

7F Posterior

El uso de ángulos de haz que atraviesan un OAR puede ser contrario a la intuición, pero es una opción.

Permite que el optimizador restrinja la dosis a los OAR con más bloqueo del MLC

NO use campos opuestos. Los campos no deben entrar y salir de la misma dirección para evitar campos POP.

IMRT - Ángulos de campo

Fields	Dose Prescription	Field Alignments	Plan Objectives	Optimization Objectives	Dose Statistics	Calculation Models	Plan Sum					
View	DVH Line	Structure	Approval Status	Plan	Course	Volume [cm ³]	Dose Cover.[%]	Sampling Cover.[%]	Min Dose [%]	Max Dose [%]	Mean Dose [%]	
☒	■	RT FEMORAL HEAD	Unapproved	7F Ant	Notes	46.8	100.0	100.0	14.8	56.7	41.8	
☒	▲	RT FEMORAL HEAD	Unapproved	7F Post	Notes	46.8	100.0	100.0	35.5	70.9	46.1	
☒	■	BOWEL	Unapproved	7F Ant	Notes	187.1	100.0	100.0	12.8	105.8	56.3	
☒	▲	BOWEL	Unapproved	7F Post	Notes	187.1	100.0	100.0	9.8	105.4	57.2	
☒	■	BLADDER	Unapproved	7F Ant	Notes	221.2	100.0	100.0	38.6	106.5	76.3	
☒	▲	BLADDER	Unapproved	7F Post	Notes	221.2	100.0	100.0	38.3	105.5	75.4	

Overview of Planning Steps

- Seleccione Arcos
- Arranque y parada de gantry
- Tamaños de campo
- Colimador
- Objetivos del plan
- Optimizar
- Revisar los resultados de optimización
- Continúe optimizando hasta que se cumplan las restricciones del plan y no haya puntos calientes fuera del PTV

Factor de modulación

IMPORTANTE: Una mayor modulación significa más flexibilidad en la planificación a expensas de más MUs y mayor tiempo de tratamiento.

Elección de los campos

- Máximo 10 arcos (típicamente 1-3)
- Número máximo de grados-1000° (aprox. 3 arcos completos)
- Longitud del arco más corto-30° grados
- Sectores de evasión (áreas en las que la tasa de dosis es de 0 mu/min), 2 por arco, longitud mínima 15°.
- El colimador no puede ser de 0° debido a fugas entre las hojas.
- Varian recomienda 45°, usamos 30°.

Two avoidance sectors per field are permitted. Minimum avoidance sector length is 15 degrees.

The beam-on sector between avoidance sectors must be at least 15 degrees, as well as the sector between an avoidance sector and the start or end of the arc.

Parámetros del haz VMAT

- El ángulo del colimador no se pone de 0° para evitar la superposición de la transmisión MLC durante el arco.
- Gire el colimador al ángulo que mejor permita que el MLC se adapte a la forma del objetivo, evite grandes rotaciones

¿Por qué arcos múltiples?

- Dispersa la modulación a varios haces
- Tal vez se necesite si el volumen entero no encaja
- Ayuda si el objetivo es tratado a niveles de dosis múltiples
- Segundo campo con colimador inverso para aumentar la modulación

Arreglos típicos

- Senos: Dos arcos de poco más de 180 grados
- Pelvis: Dos arcos completos que evitan la posible entrada a través de las barras (~300°)
- Cabeza y Cuello: Dos arcos de casi 360
- Los planes que requieren niveles de dosis múltiples o que tienen restricciones de planificación estrictas pueden requerir 3 arcos
- Cuidado con colisiones

Why is the couch important?

- CT couch different than linac
- Beam passes through the couch
- Linac couches not transparent to radiation
- Depends on the couch

Create Couch Structures

Select couch profile

Exact Couch Top with Flat panel

Movable structural rails

Left Rail Right Rail

Out In In Out

Left and right are as seen when looking towards the gantry.

CT values

Panel Surface	-450	HU
Panel Interior	-875	HU
Movable structural rails	200	HU

OK Cancel

VMAT vs. IMRT for Head and Neck

VMAT vs. IMRT para Cabeza y Cuello

Static Field IMRT: **1260 MU**

Squares: Static Field
Triangles: VMAT

VMAT: **588 MU**

la calidad de los planes generalmente similares
 Tiempo de tratamiento reducido sobre IMRT con
 campos estáticos
 MU reducido por VMAT en comparación con otras
 formas de IMRT
 El modo FFF reducirá los tiempos aún más

Los estudios NO concluyen que el VMAT resulta en una
 mejor distribución de las dosis sobre otras formas de IMRT,
 pero el VMAT ha mejorado enormemente la eficiencia de la
 administración.

VMAT a McGill

- Desde 2009
- Actualmente en 6 TrueBeams
- 65% de nuestros planes y tratamientos son con VMAT, 1% IMRT estática (técnicas heredadas)
- Todos los pacientes de VMAT en intervalos de 15 minutos
- Todos los pacientes son planificados y tratados de la misma manera

cuidados

- Piel
- Colisiones
- Sectores de evasión
- Transmisión
- La técnica no cambia la práctica
- Por ejemplo, IGRT vs VMAT vs PTV definición

¿Cómo sabemos que la dosis se administra con precisión?

- Medidas
- Cálculos

Medición de control de calidad

Total Accumulated Dose					
Target Name	Goal(Gy)	Vol Below Goal(%)	Min(Gy)	Max(Gy)	Mean(Gy)
CHAMBER	2.40		2.33	2.56	2.45
5.2mm	2.40		2.31	2.65	2.44
10.39mm	2.40		2.27	2.76	2.44
2.23mm	2.40		2.31	2.61	2.45

MLC QA

HARDWARE TESTS

120 leaf MLC

20 pairs of 0.5 cm wide leaves

40 pairs of 1 cm wide leaves

Test for leaf speed and acceleration

Test for position

Control de calidad específico del paciente: ¿por que?

- Verificar el cálculo de UM
- Verificar la entrega compleja
- Puede ser legalmente requerido
- Puede ser laborioso y consumir mucho tiempo
- ¡Compruebe lo bueno que es usted a la hora de medir las dosis!

Control de calidad específico del paciente

Control de calidad específico del paciente - película

MU checker

Point Name: Verification Beam: Subfield 1 of '1.1 RPO 205' MU: 113.00 Plan MU: 117.52 % Diff: -3.9%

Inhomogeneity Corrections Calculation Mode: SSD

Prescriptions Photon Beams MLC Data Points & Off Axis Assistance Block Entry

General

Total Calculated Dose (Gy)	1.5589
RTP Calculated Dose (Gy)	1.4933
Percent Difference (%)	4.4%
Normalize to Dose Per Fraction?	No

3D Coordinates

Enter 3D Coordinates for Points?

X Position: -0.04 cm Y Position: -0.04 cm Z Position: 0.00 cm

Beam Desc.	Depth	Equiv. Path	SSD	OADx	OADz	Dose	RTP Dose	% Diff.
1 Subfield 0 of '1.1 RPO 205'	14.90	N/A	85.10	0.00	0.00	0.0094	0.0066	43.1%
2 Subfield 1 of '1.1 RPO 205'	14.90	N/A	85.10	0.00	0.00	0.2237	0.2150	4.0%
3 Subfield 2 of '1.1 RPO 205'	14.90	N/A	85.10	0.00	0.00	0.0091	0.0066	37.9%
4 Subfield 0 of '1.2 RPO 255'	14.90	N/A	85.10	0.00	0.00	0.0065	0.0044	47.2%
5 Subfield 1 of '1.2 RPO 255'	14.90	N/A	85.10	0.00	0.00	0.2305	0.2262	1.9%
6 Subfield 0 of '1.3 RAO 310'	14.90	N/A	85.10	0.00	0.00	0.0210	0.0198	6.3%
7 Subfield 1 of '1.3 RAO 310'	14.90	N/A	85.10	0.00	0.00	0.2147	0.2063	4.0%
8 Subfield 0 of '1.4 ANT 0'	15.00	N/A	85.00	0.00	0.00	0.0090	0.0066	36.1%
9 Subfield 1 of '1.4 ANT 0'	15.00	N/A	85.00	0.00	0.00	0.1284	0.1251	2.6%
10 Subfield 2 of '1.4 ANT 0'	15.00	N/A	85.00	0.00	0.00	0.0090	0.0066	36.7%
11 Subfield 0 of '1.5 LAO 50'	14.90	N/A	85.10	0.00	0.00	0.2409	0.2306	4.5%
12 Subfield 1 of '1.5 LAO 50'	14.90	N/A	85.10	0.00	0.00	0.0127	0.0110	14.8%
13 Subfield 0 of '1.6 LPO 105'	14.90	N/A	85.10	0.00	0.00	0.2445	0.2395	2.1%

Display

Mag Factor: 1.00

Toward Gantry ↑

↓ Away From Gantry

Display

Mag Factor: 1.00

The currently selected beam does not have a valid intensity map. The doses will not automatically be updated. The intensity map can be re-computed by pressing either the "Compute Beam" or "Compute All Beams" button on the toolbar.

QA results table

YEAR	TPS	#QA	Measured/ calc	% from calc	ST DEV	CI (95%)
2002	CORVUS	41	0.985	-1.488	0.041	9.6
2003	CORVUS	87	0.972	-2.839	0.036	10
2004	CORVUS	125	0.998	-0.174	0.024	4.9
2005	CORVUS	134	1.005	0.457	0.025	5.4
2006	CORVUS	182	0.998	-0.238	0.021	4.4
2007	CORVUS	164	0.984	-1.578	0.019	5.3
2008	CORVUS	223	0.988	-1.239	0.024	6.0
2009	ECLIPSE	25	1.010	1.032	0.014	3.8
2009 (RA)	ECLIPSE	98	1.013	1.341	0.015	4.2
RADCALC	ECLIPSE	205	1.022	2.184	0.012	4.6

Tomotherapy DQA

All Tomotherapy DQA - McGill University

N=486 (March 2007 - October 2009)

All Tomotherapy DQA - McGill University

N=486 (March 2007 - October 2009)

PSQAs – VMAT 2009

McGill action levels for IMRT QA

QA method	Measurement point	Action level	Action if level exceeded
Software (RadCalc)	High dose point	$\pm 5\%$	Perform IC/film mapcheck
Ion chamber	High dose point	$\pm 5\%$	Redo QA
Film/mapcheck	High isodose lines	3 mm DTA	Redo QA
filmFilm/mapcheck	Isodose distribution	90% pixels pass 5%, 3mm	Investigate/rationalize
Audit of point measurements	High dose point	90% of points $\pm 3\%$ per year	Investigate/rationalize

PSQAs – VMAT 2009

107 VMAT Mapcheck QA 2009-2010 3%, 3mm

VMAT Puesta en marcha 2015

Done Both IC & M	Plan	Plan Type (RARC MRT)	Machine	Ion Chamber QA	Physicist	homogeneous IC calc	Mapcheck	Physicist2	RADCALC %DIFF TO CALC POINT	MapCheck port attached	Site	1 MU	2 MU	3 MU	Column5	Total dose	fractions	dose per f
1	FP2 PROSTATE	RARC	TB3	-0.27	TC	HEToff	98.1	ks	1.60%		PROSTATE	246	252			34.2	19	1.8
2	GP1 BRAIN	RARC	STX1	0.97	GA	HEToff	97.6	Ks	0.50%		BRAIN	204	213			52.2	29	1.8
3	FP1 ABDO	RARC	TB6	-0.18	TC	HEToff	98.8	IA	0.60%		ABDOMEN	453	320	245		50.4	28	1.8
4	FP2 BREAST	RARC	TB4	-1.49	TC	HEToff	100	KS	0.80%		BREAST	223	241			10	4	2.5
5	FP1 ENT	RARC	TB6	0.376	NS/WP	HEToff	97.7	GA/IA	2.70%		ENT	510	446			70	35	2
6	FP1 LUNG	RARC	TB6	0.23	TC	HEToff	98.5	IA	-0.80%		LUNG	376	471			56	28	2
7	FP2 LT BREAST	RARC	TB4	-0.22	TC	HEToff	95.9	GA/IA	3.30%		BREAST	399	409			12.5	5	2.5
8	GP1 PROSTATE	RARC	TB6	-0.19	TC	HEToff	98.7	IA	1.40%		PROSTATE	367	383			50	20	2.5
9	FP1 PELVIS	RARC	STX1	-2.07	GA	HEToff	99.3	Ks	1.30%		RECTUM	491	501			25	5	5
11	M1P1_ENT	RARC	TB6	0.4	TC	HEToff	98.7	GA/IA			ENT	179	175			66	30	2.2
12	FP2 RT BREAST	RARC	TB5	3.67	TC	HEToff	96.6	GA/IA	-0.10%		BREAST	388	332			12.5	5	2.5
13	GP1 RECTUM	RARC	TB3	-0.5	GA	HEToff	97.7	Ks	-0.80%		RECTUM	285	298			45	25	1.8
14	FP2 RECTUM	RARC	TB3	-0.2	GA	HEToff	95.6	ks	3.00%		RECTUM	263	283			5.4	3	1.8
15	GP1_BRAIN	RARC	STX1	0.05	NS	HEToff	98.8	NS	1.90%		BRAIN	209	228			50	25	2
16	FP2_BRAIN	RARC	STX1	-0.04	NS	HEToff	99	NS	2.80%		BRAIN	173	226			10	5	2
17	FP1 RT LUNG	RARC	TB6	-1.04	TC	HEToff	99.1	IA	1.80%		LUNG	257	295			60	30	2
18	GP1_RECTUM	RARC	TB6	-1.356	NS/WP	HEToff	93.6	GA/IA	-2.10%		RECTUM	197	204			45	25	1.8
19	FP2_RECTUM	RARC	TB6	-0.795	NS/WP	HEToff	94.9	GA/IA	-1.00%		RECTUM	234	250			5.4	3	1.8
20	FP2_PROSTATE	RARC	TB6	-1.05	TC	HEToff	92.4	IA	1.30%		PROSTATE	319	343			46	23	2
21	FP1 PELVIS	RARC	TB6	-0.6	TC	HEToff	98.8	GA/IA	1.50%	YES	PELVIS	276	314			50	25	2
22	GP1 RECTUM	RARC	TB4	-1.27	TC	HEToff	98.2	Ks	-0.20%		RECTUM	228	242			45	25	1.8
23	FP2 RECTUM	RARC	TB4	-1.22	TC	HEToff	98.8	Ks	1.70%		RECTUM	181	201			9	5	1.8
24	FP1 SPINE	RARC	STX2	1.5	GA	HEToff	98.5	GA/IA	2.30%		SPINE	839	822	846		18	2	9
25	FP2_RT BREAST	RARC	TB4	0.046	TC	HEToff	100	Ks	0.70%		BREAST	200	195			10	4	2.5
26	GP1_RT LEG	RARC	TB4	-1.1	TC	HEToff	99.4	Ks	-0.40%		RT LEG	179	247			50	25	2
27	FP1_PROSTATE	RARC	TB6	0.02	TC	HEToff	97.1	IA	1.30%		PROSTATE	292	376			44	22	2
28	FP2_PROSTATE	RARC	TB6	-0.24	TC	HEToff	96.4	IA	1.80%		PROSTATE	352	355			22	11	2
29	FP2 LT BREAST	RARC	TB3	-1.2	GA	HEToff	98.1	GA/IA	0.00%		BREAST	302	309			10	4	2.5
30	FP1 BRAIN	RARC	STX1	-1.64	GA	HEToff	99.2	Ks	3.50%		BRAIN	387	353			20	5	4
31	FP1_RT LEG	RARC	TB3	-1.4	GA	HEToff	98.7	Ks	0.50%		LEG	225	200			50	25	2
32	FP1 NECK	RARC	TB6	0.25	TC	HEToff	98.8	IA	1.00%		ENT	366	372			4	2	2
33	GP1 ENT	RARC	STX1	0.388	GA	HEToff	97.2	GA/IA	4.90%		ENT	211	247			70	35	2
34	FP2_LT BREAST	RARC	TB4	-0.87	TC	HEToff	99.1	Ks	0.00%		BREAST	379	380			10	4	2.5
34	GP1 ENT	RARC	TB5	2.87	TC	HEToff	99	Ks	3.40%		ENT	303	346			70	35	2
36	GP1 PELVIS	RARC	TB5	-2.01	TC	HEToff	98.3	IA	0.70%		PELVIS	265	258			30.6	17	1.8
37	FP1_ENT	RARC	TB5	0.1	TC	HEToff	100	GA/IA	1.30%	YES	ENT	563	524			35	10	3.5
38	FP2 PELVIS	RARC	TB5	-1.45	TC	HEToff	92.5	IA	1.80%		PELVIS	203	207			19.8	11	1.8
39	FP1 PROSTATE	RARC	STX1	0.998	GA	HEToff	95.4	Ks	0.40%		PROSTATE	213	160			79.2	44	1.8
40	GP1 RT LUNG	RARC	TB6	0.51	TC	HEToff	99.2	IA	1.10%		LUNG	414	390			45	25	1.8
41	FP1 PROSTATE	RARC	TB3	0.91	TC	HEToff	99.6	Ks	0.70%		PROSTATE	338	327			62.5	25	2.5
42	GP1 BLADDER	RARC	STX1	-6.413	GA	HEToff	99.9	GA/IA	0.60%		BLADDER	240	236			50	20	2.5
43	FP1_ENT	RARC	TB5	0.19	TC	HEToff	99.8	IA	1.70%		ENT	234	224			70	35	2
44	FP1 BLADDER	RARC	TB6	-0.05	TC	HEToff	99.7	IA	-0.50%		BLADDER	490	771			20	4	5
45	FP1 PELVIS	RARC	TB6	-0.24	TC	HEToff	98.9	IA	1.40%		PELVIS	318	323					2
46	FP1_LT THIGH	RARC	TB4	-1.43	TC	HEToff	99.7	Ks	0.50%		THIGH	221	227					2
47	FP2_PROSTATE	RARC	TB6	-1.1	TC	HEToff	96.9	IA	1.30%		PROSTATE	234	314			22	11	2

Screenshot

VMAT Puesta en marcha 100 PSQA

Verificando el verificador MU!

**Report of
IMRT Head and Neck
Phantom Irradiation**

Date of Report: May 28, 2015
Institution: McGill University Health Centre
Physician: Russell Ruo
Radiation Machine: Varian, TrueBeam (1321) - 6 MV
Intensity Modulation Device: Multileaf Collimator
IMRT Technique: Rapid Arc
Treatment Planning System: Varian, Eclipse (IMRT) - AAA
Date of Irradiation: May 1, 2015

(Actually STX1
did not have right
serial #?)

Description of Procedure

An anthropomorphic head phantom incorporating a rectangular dosimetry insert was imaged and irradiated to approximately 6.6 Gy using an IMRT technique. The dosimetry insert consisted of one primary PTV containing four TLD capsules, a secondary PTV and an organ at risk (OAR), each containing two TLD capsules. The TLD capsules provided point dose information. Three sheets of GAF Chromic™ Dosimetry Media provided dose profiles through the center of primary PTV.

The dosimetric precision of the TLD is 3%, and the spatial precision of the film and densitometer system is 1 mm.

Summary of TLD and film results:

Location	IROC-H vs. Inst.	Criteria	Acceptable
Primary PTV sup. ant.	1.00	0.93 - 1.07	Yes
Primary PTV inf. ant.	1.01	0.93 - 1.07	Yes
Primary PTV sup. post.	1.01	0.93 - 1.07	Yes
Primary PTV inf. post.	1.01	0.93 - 1.07	Yes
Secondary PTV sup.	1.00	0.93 - 1.07	Yes
Secondary PTV inf.	1.03	0.93 - 1.07	Yes

Film Plane	Gamma Index*	Criteria	Acceptable
Axial	100%	≥85%	Yes
Sagittal	99%	≥85%	Yes

*Percentage of points meeting gamma-index criteria of 7% and 4 mm.

The phantom irradiation results listed in the table above **do meet** the criteria established by IROC Houston in collaboration with the cooperative study groups. Therefore, your institution **has satisfied** the phantom irradiation component of the credentialing process to enter patients into certain protocols that allow the use of IMRT.

TLD and Film Analysis by: Nadia Hernandez and Andrea Molineu, M.S.

Report Checked by:

David S. Folowill
David S. Folowill, Ph.D.
Director, IROC Houston QA Center

Isodoses (Gy)

6.90
6.60
6.27
5.40
5.10
4.50
3.00
1.50

Conclusiones VMAT

- tratamiento rápido
- distribución conforme
- puede beneficiar al 90% de los pacientes
- “workflow” único para tecnólogos

Cone 4 mm

Cone 40 mm

80 - 50% : 2 mm

80 - 20% : 5 mm

80 - 50% : 6 mm

80 - 20% : 16 mm